



# Reston Hospital Center

*HCA Virginia Health System*

An HCA affiliate

# Ann B. Rodriguez Cancer Center


*Helping cancer patients along their  
journey from diagnosis to  
survivorship...*



# Cancer Program and Goals

## *Passion for Excellence, Compassion for You*


- Providing exceptional care in a timely manner
- Seamless coordination of services for optimal outcomes
- Personalized high level of service
- Highest level of excellence within community
- Multidisciplinary approach to patient care
- Evidence based quality patient care
- Access to clinical trials



# 2010-2013 Total Cancer Cases –Top Sites


**Total Cases**  
**2010=679**  
**2011=743**  
**2012=721**  
**2013=889**

■ 2013  
■ 2012  
■ 2011  
■ 2010




# 2010-2013 Total Cancer Cases

## Cancer Cases by Stage



## Analytic Cancer Cases






# Quality Measures and Outcomes

Cancer Program Practice Profile Reports for Breast, Colon, and Rectal Cancer		2008	2009	2010	2011
<b>Breast</b>	Radiation therapy is administered within 1 year if diagnosis for women under age 70 receiving breast conserving surgery for breast cancer	100%	98%	96.5%	93.4%
<b>Breast</b>	Combination chemotherapy is considered or administered within 4 months of diagnosis for women under 70 with AJCC T1c NO MO, of Stage II or III ERA and or PRA positive breast cancer	90%	100%	90%	85.7%
<b>Breast</b>	Tamoxifen or third generation aromatase inhibitors is considered or administered within 1 year of diagnosis for women with AJCC T1c No MO of Stage II or III ERA and or PRA positive breast cancer.	98.3%	97.8%	100%	95.9%
<b>Colon</b>	Adjuvant chemotherapy is considered or administered within 4 months (120 days) of diagnosis for patients under the age of 80 with AJCC Stage III (lymph node positive) colon cancer. <b>[ACT]</b>	100%	100%	100%	100%
<b>Colon</b>	At least 12 regional lymph nodes are removed and pathologically examined for resected colon cancer. <b>[12RLN]</b>	93.3%	85.2%	85.7%	100%
<b>Rectal</b>	Radiation therapy is considered or administered within 6 months (180 days) of diagnosis for patients under the age of 80 of with clinical or pathologic AJCC T4N0M0 or Stage III receiving surgical resection for rectal cancer. <b>[AdjRT]</b>	100%	100%	88.9%	100%

# Unadjusted 5 Year Breast Cancer Survival Rates by Stage, 2003 - 2006



	Stage 0	Stage I	Stage II	Stage III	Stage IV	OVERALL
<b>My Facility</b>	99.2 %	97.7 %	91.5 %	79.1 %	*Too few cases	93.2 %
<b>All CoC</b>	95.6 %	92.2 %	85.4 %	66.7 %	21.1 %	85.5 %

DISCLAIMER: (Includes patients of all ages, deaths from all causes, and there has been no adjustment for co-morbidities or other risk factors. If no result is shown, too few cases were submitted). Survival not calculated if less than 30 cases